

INSTRUCTIONS FOR FLEXIBLE SIGMOIDOSCOPY

If you have a history of constipation (move your bowels 2-3 times weekly) you may need additional bowel prep to help clear your colon. Please contact the intake nurse to discuss prep options.

STEP 1 7 DAYS BEFORE PROCEDURE

1

- Stop taking Phentermine (Adipex)
- Avoid foods that contain Olestra/Oean (ex: fat free potato chips).
- Avoid foods that have seeds, nuts, lettuce, cabbage, corn and beans.
- Stop fiber supplements.
- Purchase at pharmacy: (Supplies may be purchased without a prescription)
Ducolax suppository
Fleet Enema (2 needed)
- If taking Plavix, Effient, Aggrenox, Coumadin, Eliquis, Pradaxa or any blood thinning medication, please inform our nurse at the time of call. This medication will need to be held prior to the procedure and an order obtained from the prescribing physician.

If diabetic, please contact your primary care doctor regarding instructions for diabetic medication while on the clear liquid diet.

STEP 2 DAY BEFORE PROCEDURE

2

- Insert one Dulcolax suppository rectally at 6:00 p.m. on the night before the exam.

STEP 3 DAY OF PROCEDURE

3

- Two hours before your appointment time on the day of your procedure, take one Fleet enema. Hold for five or ten minutes, then expel.
- After expelling first enema, repeat with a second Fleet enema.
- You may continue your usual diet with this preparation, unless your preparation is scheduled with **sedation**. If your procedure is scheduled with sedation you are to have **nothing by mouth eight hours prior** to your procedure.

STEP 4 8 HOURS BEFORE PROCEDURE DO NOT DRINK ANY FURTHER FLUIDS.

4

- 2 hours before your procedure, take heart and blood pressure medication with a small sip of water.
- Please do not apply lotions, perfumes or cologne the morning of your procedure.
- **Females may be required to do a urine pregnancy test prior to the procedure. Please inquire (about test) at the desk upon arrival or before using the restroom.**

You should take your heart medications (excluding Plavix), blood pressure medications, and medications for lung problems on the day of the exam with a sip of water. All other medications should not be taken until after the procedure

You **may** continue taking aspirin and aspirin products unless specifically instructed not to by your physician. Do **not** stop taking prednisone, methotrexate or Enbrel.

If you are unsure about taking any medication in conjunction with your procedure, please do not hesitate to call our Intake nurse at (309)370-6838.

You must have someone drive you home if having sedation. You may not go home by bus or taxicab

TIPS FOR IMPROVED BOWEL PREP: TO AVOID A SORE BOTTOM FROM FREQUENT BOWEL MOVEMENTS:

- Apply diaper rash cream/zinc oxide LIBERALLY after each trip to the bathroom
- Avoid rubbing; gently pat area with wet washcloth or pre-moistened wipe or rinse with water
- Wipe with refrigerated Tucks or Fleet relief pads (available over the counter)
- Have reading material handy in the bathroom, MP3 player, iPod, etc.